

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
6 December 2001 (06.12.2001)

PCT

(10) International Publication Number  
**WO 01/92545 A1**

(51) International Patent Classification<sup>7</sup>: C12N 15/70,  
15/77, C12P 13/08, C12N 1/21 // 15/31

(21) International Application Number: PCT/EP01/03980

(22) International Filing Date: 6 April 2001 (06.04.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
100 26 494.8 27 May 2000 (27.05.2000) DE  
101 02 823.7 23 January 2001 (23.01.2001) DE

(71) Applicant: DEGUSSA AG [DE/DE]; Bennigsenplatz 1,  
40474 Düsseldorf (DE).

(72) Inventor: RIEPING, Mechthild; Mönkebergstrasse 1,  
33619 Bielefeld (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,

DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: PROCESS FOR THE FERMENTATIVE PREPARATION OF L-THREONINE



(57) Abstract: The invention provides a process for the fermentative preparation of L-threonine using Enterobacteriaceae which in particular already produce L-threonine and in which the nucleotide sequence(s) of coryneform bacteria which code(s) for the thrE gene are enhanced, in particular over-expressed.

**Process for the fermentative preparation of L-threonine**

This invention relates to a process for the fermentative preparation of L-threonine using Enterobacteriaceae in which the thrE gene of coryneform bacteria is enhanced.

**5 Prior art**

L-Threonine is used in animal nutrition, in human medicine and in the pharmaceuticals industry. It is known that L-threonine can be prepared by fermentation of strains of Enterobacteriaceae, in particular Escherichia coli and  
10 Serratia marcescens. Because of their great importance, work is constantly being undertaken to improve the preparation processes. Improvements to the process can relate to fermentation measures, such as e.g. stirring and supply of oxygen, or the composition of the nutrient media,  
15 such as e.g. the sugar concentration during the fermentation, or the working up to the product form, by e.g. ion exchange chromatography, or the intrinsic output properties of the microorganism itself.

Methods of mutagenesis, selection and mutant selection are  
20 used to improve the output properties of these microorganisms. Strains which are resistant to antimetabolites, such as e.g. the threonine analogue  $\alpha$ -amino- $\beta$ -hydroxyvaleric acid (AHV), or are auxotrophic for amino acids of regulatory importance and produce L-  
25 threonine are obtained in this manner.

Methods of the recombinant DNA technique have also been employed for some years for improving the strain of Enterobacteriaceae strains which produce L-threonine, by amplifying individual threonine biosynthesis genes and  
30 investigating the effect on the L-threonine production.

## Object of the invention

The inventors had the object of providing new measures for improved fermentative preparation of L-threonine.

## Description of the invention

- 5 The invention provides a process for the fermentative preparation of L-threonine using Enterobacteriaceae which in particular already produce L-threonine and in which the nucleotide sequence(s) of coryneform bacteria which code(s) for the thrE gene are enhanced, in particular over-  
10 expressed.

In particular, the process is a process for the preparation of L-threonine, which comprises carrying out the following steps:

- 15 a) fermentation of microorganisms of the family Enterobacteriaceae in which at least the thrE gene of coryneform bacteria is enhanced (over-expressed), optionally in combination with further genes,
- b) concentration of the L-threonine in the medium or in the cells of the microorganisms of the family  
20 Enterobacteriaceae, and
- c) isolation of the L-threonine.

The term "enhancement" in this connection describes the increase in the intracellular activity of one or more enzymes or proteins in a microorganism which are coded by  
25 the corresponding DNA, for example by increasing the number of copies of the gene or genes, using a potent promoter or a gene which codes for a corresponding enzyme or protein with a high activity, and optionally combining these measures.

- 30 The microorganisms which the present invention provides can prepare L-threonine from glucose, sucrose, lactose, fructose, maltose, molasses, starch, cellulose or from

glycerol and ethanol. They are representatives of Enterobacteriaceae, in particular of the genera Escherichia and Serratia. Of the genus Escherichia the species Escherichia coli and of the genus Serratia the species

5 Serratia marcescens are to be mentioned in particular.

Suitable L-threonine-producing strains of the genus Escherichia, in particular of the species Escherichia coli, are, for example

Escherichia coli TF427  
10 Escherichia coli H4578  
Escherichia coli KY10935  
Escherichia coli VNIIgenetika MG-442  
Escherichia coli VNIIgenetika M1  
Escherichia coli VNIIgenetika 472T23  
15 Escherichia coli BKIIM B-3996  
Escherichia coli kat 13  
Escherichia coli KCCM-10132

Suitable L-threonine-producing strains of the genus Serratia, in particular of the species Serratia marcescens,

20 are, for example

Serratia marcescens HNr21  
Serratia marcescens TLR156  
Serratia marcescens T2000

It has been found that Enterobacteriaceae produce L-

25 threonine in an improved manner after over-expression of the thrE gene of coryneform bacteria which codes for threonine export.

Nucleotide sequences of the thrE gene of coryneform bacteria are shown in SEQ ID No 1 and SEQ ID No 3 and the

30 resulting amino acid sequences of the export proteins are shown in SEQ ID No 2 and SEQ ID No 4.

The thrE gene shown in SEQ ID No 1 and SEQ ID No 3 can be used according to the invention. Alleles of the thrE gene of coryneform bacteria which result from the degeneracy of

the genetic code or due to sense mutations of neutral function can furthermore be used.

To achieve an over-expression, the number of copies of the corresponding genes can be increased, or the promoter and regulation region or the ribosome binding site upstream of the structural gene can be mutated. Expression cassettes which are incorporated upstream of the structural gene act in the same way. By inducible promoters, it is additionally possible to increase the expression in the course of fermentative L-threonine production. The expression is likewise improved by measures to prolong the life of the m-RNA. Furthermore, the enzyme activity is also increased by preventing the degradation of the enzyme protein. The genes or gene constructions can either be present in plasmids with a varying number of copies, or can be integrated and amplified in the chromosome. Alternatively, an over-expression of the genes in question can furthermore be achieved by changing the composition of the media and the culture procedure.

Instructions in this context can be found by the expert, inter alia, in Chang and Cohen (Journal of Bacteriology 134:1141-1156 (1978)), in Hartley and Gregori (Gene 13:347-353 (1981)), in Amann and Brosius (Gene 40:183-190 (1985)), in de Broer et al. (Proceedings of the National of Sciences of the United States of America 80:21-25 (1983)), in LaVallie et al. (BIO/TECHNOLOGY 11, 187-193 (1993)), in PCT/US97/13359, in Llosa et al. (Plasmid 26:222-224 (1991)), in Quandt and Klipp (Gene 80:161-169 (1989)), in Hamilton (Journal of Bacteriology 171:4617-4622 (1989)), in Jensen and Hammer (Biotechnology and Bioengineering 58, 191-195 (1998) and in known textbooks of genetics and molecular biology.

Plasmid vectors which are capable of replication in Enterobacteriaceae, such as e.g. cloning vectors derived from pACYC184 (Bartolomé et al.; Gene 102, 75-78 (1991)), pTrc99A (Amann et al.; Gene 10:211-216 (1981)), or p1111

derivatives (Vocke and Bastia, Proceedings of the National Academy of Science USA 80 (21):6557-6561 (1983)) can be used. A strain transformed with a plasmid vector where the plasmid vector carries the nucleotide sequence which codes  
5 for the thrE gene of coryneform bacteria can be employed in a process according to the invention.

In addition, it may be advantageous for the production of L-threonine with strains of the family Enterobacteriaceae to over-express one or more enzymes of the known threonine  
10 biosynthesis pathway or enzymes of anaplerotic metabolism or enzymes for the production of reduced nicotinamide adenine dinucleotide phosphate, in addition to the thrE gene of coryneform bacteria. Thus, for example

- at the same time the thrABC operon which codes for  
15 aspartate kinase, homoserine dehydrogenase, homoserine kinase and threonine synthase (US-A-4,278,765), or
- at the same time the pyc gene which codes for pyruvate carboxylase (DE-A-19 831 609), or
- at the same time the pps gene which codes for  
20 phosphoenol pyruvate synthase (Molecular and General Genetics 231:332 (1992)), or
- at the same time the ppc gene which codes for phosphoenol pyruvate carboxylase (Gene 31:279-283 (1984)), or
- 25 • at the same time the genes pntA and pntB which code for transhydrogenase (European Journal of Biochemistry 158:647-653 (1986)), or
- at the same time the gdhA gene which codes for glutamate dehydrogenase (Gene 27:193-199 (1984)), or
- 30 • at the same time the rhtB gene which imparts L-homoserine resistance (EP-A-0994190)

can be obtained in particular from the following:

In addition to over-expression of the thrE gene it may furthermore be advantageous, for the production of L-threonine, to eliminate undesirable side reactions, such as e.g. threonine dehydrogenase (Nakayama: "Breeding of Amino  
5 Acid Producing Microorganisms", in: Overproduction of Microbial Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982 and Bell and Turner, Biochemical Journal 156, 449-458 (1976)). Bacteria in  
10 L-threonine are at least partly eliminated can be employed in the process according to the invention.

The microorganisms produced according to the invention can be cultured in the batch process (batch culture) or in the fed batch process (feed process). A summary of known  
15 culture methods is described in the textbook by Chmiel (Bioprozesstechnik 1. Einführung in die Bioverfahrenstechnik [Bioprocess Technology 1. Introduction to Bioprocess Technology (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und  
20 periphere Einrichtungen [Bioreactors and Peripheral Equipment] (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

The culture medium to be used must meet the requirements of the particular strains in a suitable manner. Descriptions of culture media for various microorganisms are contained  
25 in the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981). Sugars and carbohydrates, such as e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and cellulose, oils and fats, such as e.g.  
30 soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, such as e.g. palmitic acid, stearic acid and linoleic acid, alcohols, such as e.g. glycerol and ethanol, and organic acids, such as e.g. acetic acid, can be used as the source of carbon. These substance can be used  
35 individually or as a mixture. Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea,

or inorganic compounds, such as ammonium sulphate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate, can be used as the source of nitrogen. The sources of nitrogen can be used individually or as a  
5 mixture. Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts can be used as the source of phosphorus. The culture medium must furthermore comprise salts of metals, such as e.g. magnesium sulfate or iron  
10 sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be employed in addition to the abovementioned substances. Suitable precursors can moreover be added to the culture medium. The starting substances mentioned can  
15 be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.

Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia or aqueous ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed  
20 in a suitable manner to control the pH. Antifoams, such as e.g. fatty acid polyglycol esters, can be employed to control the development of foam. Suitable substances having a selective action, e.g. antibiotics, can be added to the medium to maintain the stability of plasmids. To  
25 maintain aerobic conditions, oxygen or oxygen-containing gas mixtures, such as e.g. air, are introduced into the culture. The temperature of the culture is usually 25°C to 45°C, and preferably 30°C to 40°C. Culturing is continued until a maximum of L-threonine has formed. This target is  
30 usually reached within 10 hours to 160 hours.

The analysis of L-threonine can be carried out by anion exchange chromatography with subsequent ninhydrin derivatization, as described by Spackman et al. (Analytical Chemistry, 30, (1958), 1190), or it can take place by  
35 reversed phase HPLC as described by Lindroth et al. (Analytical Chemistry (1979) 51: 1167-1174).



The following microorganism has been deposited at the Deutsche Sammlung für Mikroorganismen und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) in accordance with the Budapest

5 Treaty:

- *Brevibacterium flavum* strain DM368-2 pZ1thrE as DSM 12840

The plasmid pZ1thrE contains the thrE gene of *Corynebacterium glutamicum* ATCC13032.

- 10 The present invention is explained in more detail in the following with the aid of embodiment examples.

The isolation of plasmid DNA from *Escherichia coli* and all techniques of restriction, Klenow and alkaline phosphatase treatment were carried out by the method of Sambrook et al.

- 15 (Molecular cloning - A laboratory manual (1989) Cold Spring Harbour Laboratory Press). The transformation of *Escherichia coli* was carried out by the method of Chung et al. (Proceedings of the National Academy of Sciences of the United States of America USA (1989) 86: 2172-2175).

- 20 The incubation temperature during the preparation of *E. coli* strains and transformants was 37°C. In the gene replacement process of Hamilton et al. (Journal of Bacteriology (1989) 171: 4617-4622) temperatures of 30°C and 44°C were used.

25 Example 1

Cloning and sequencing of the thrE gene of *Corynebacterium glutamicum* ATCC14752

1. Transposon mutagenesis and mutant selection

- The strain *Corynebacterium glutamicum* ATCC14752ΔilvA was  
30 subjected to mutagenesis with the transposon Tn5531, the sequence of which is deposited under accession number U53587 in the nucleotide databank of the National Center

for Biotechnology Information (Bethesda, USA). The incorporation of a deletion in the *ilvA* gene of *Corynebacterium glutamicum* ATCC14752 was carried out with the gene exchange system described by Schäfer et al. (Gene 5 (1994) 145: 69-73). For this, an inactivation vector pK19mobsacB $\Delta$ *ilvA* constructed by Sahm et al. (Applied and Environmental Microbiology (1999) 65: 1973-1979) was used for the deletion. The methylase-defective *Escherichia coli* strain SCS110 (Jerpseth and Kretz, STRATEGIES in molecular 10 biology 6, 22, (1993)) from Stratagene (Heidelberg, Germany) was first transformed with 200 ng of the vector pK19mobsacB $\Delta$ *ilvA*. Transformants were identified with the aid of their kanamycin resistance on LB agar plates containing 50  $\mu$ g/mL kanamycin. The plasmid pK19mobsacB $\Delta$ *ilvA* 15 was prepared from one of the transformants. By means of electroporation (Haynes et al., FEMS Microbiology Letters (1989) 61: 329-334), this inactivation plasmid was then introduced into the strain *Corynebacterium glutamicum* ATCC14752. Clones in which the inactivation vector was 20 present integrated in the genome were identified with the aid of their kanamycin resistance on LBHIS agar plates containing 15  $\mu$ g/mL kanamycin (Liebl et al., FEMS Microbiology Letters (1989) 65: 299-304). To select for the excision of the vector, kanamycin-resistant clones were 25 plated out on sucrose-containing LBG medium (LB medium with 15 g/L agar, 2% glucose and 10% sucrose). This gave colonies which have lost the vector again by a second recombination event (Jäger et al.; Journal of Bacteriology (1992) 174: 5462-5465). By transinoculation on to minimal 30 medium plates (CGXII medium with 15 g/L Agar (Keilhauer et al., Journal of Bacteriology (1993) 175: 5595-5603)) with and without 300 mg/L L-isoleucine, or with and without 50  $\mu$ g/mL kanamycin, six clones were isolated which were kanamycin-sensitive and isoleucine-auxotrophic due to 35 excision of the vector and in which the incomplete *ilvA* gene ( $\Delta$ *ilvA* allele) was now present in the genome. One of these clones was designated strain ATCC14752 $\Delta$ *ilvA* and employed for the transposon mutagenesis.

From the methylase-defective *E. coli* strain GM2929pCGL0040 (*E. coli* GM2929: Palmer et al., Gene (1994) 143: 1-12) was isolated the plasmid pCGL0040 (figure 1), which contains the combined transposon Tn5531 (Ankri et al., Journal of Bacteriology (1996) 178: 4412-4419). The strain *Corynebacterium glutamicum* ATCC14752 $\Delta$ ilvA was transformed by means of electroporation (Haynes et al., FEMS Microbiology Letters (1989) 61: 329-334) with the plasmid pCGL0040. Clones in which the transposon Tn5531 was integrated into the genome were identified with the aid of their kanamycin resistance on LBHIS agar plates containing 15  $\mu$ g/mL kanamycin (Liebl et al., FEMS Microbiology Letters (1989) 65: 299-304). 2000 clones were obtained in this manner, and were investigated for delayed growth in the presence of threonyl-threonyl-threonine. For this, all the clones were transferred individually to CGXII minimal medium agar plates with and without 2 mM threonyl-threonyl-threonine. The medium was identical to the CGXII medium described by Keilhauer et al. (Journal of Bacteriology (1993) 175: 5593-5603), but additionally contained 25  $\mu$ g/mL kanamycin, 300 mg/L L-isoleucine and 15 g/L agar. The composition of the medium described by Keilhauer et al. is shown in table 1.

Table 1

Composition of the CGXII medium

Component	Concentration
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	20 g/L
Urea	5 g/L
KH <sub>2</sub> PO <sub>4</sub>	1 g/L
K <sub>2</sub> HPO <sub>4</sub>	1 g/L
MgSO <sub>4</sub> x 7 H <sub>2</sub> O	0.25 g/L
3-Morpholinopropanesulfonic acid	42 g/L
CaCl <sub>2</sub>	10 mg/L
FeSO <sub>4</sub> x 7 H <sub>2</sub> O	10 mg/L
MnSO <sub>4</sub> x H <sub>2</sub> O	10 mg/L
ZnSO <sub>4</sub> x 7H <sub>2</sub> O	1 mg/L

CuSO <sub>4</sub>	0.2 mg/L
NiCl <sub>2</sub> x 6 H <sub>2</sub> O	0.02 mg/L
Biotin	0.2 mg/L
Glucose	40 g/L
Protocatechuic acid	30 mg/L

The agar plates were incubated at 30°C and the growth was investigated after 12, 18 and 24 hours. A transposon mutant was obtained which grew in a manner comparable to the starting strain *Corynebacterium glutamicum* ATCC14752ΔilvA without threonyl-threonyl-threonine but showed delayed growth in the presence of 2 mM threonyl-threonyl-threonine. This was designated ATCC14752ΔilvAthrE::Tn5531.

## 2. Cloning and sequencing of the insertion site of Tn5531 in ATCC14752ΔilvAthrE::Tn5531

To clone the insertion site lying upstream of the transposon Tn5531 in the mutant described in example 1.1, the chromosomal DNA of this mutant strain was first isolated as described by Schwarzer et al. (Bio/Technology (1990) 9: 84-87) and 400 ng thereof were cleaved with the restriction endonuclease EcoRI. The complete restriction batch was ligated in the vector pUC18 (Norander et al., Gene (1983) 26: 101-106, Roche Diagnostics, (Mannheim, Germany), which was also linearized with EcoRI. The *E. coli* strain DH5αmc<sup>r</sup> (Grant et al., Proceedings of the National Academy of Sciences of the United States of America USA (1990) 87: 4645-4649) was transformed with the entire ligation batch by means of electroporation (Dower et al., Nucleic Acid Research (1988) 16: 6127-6145). Transformants in which the insertion sites of the transposon Tn5531 were present cloned on the vector pUC18 were identified with the aid of their carbenicillin and kanamycin resistance on LB agar plates containing 50 µg/mL carbenicillin and 25 µg/mL kanamycin. The plasmids were prepared from three of the transformants and the sizes of

- the cloned inserts were determined by restriction analysis. The nucleotide sequence of the insertion site on one of the plasmids with an insert approx. 5.7 kb in size was determined by the dideoxy chain termination method of Sanger et al. (Proceedings of the National Academy of Sciences of the United States of America USA (1977) 74: 5463-5467). For this, 2.2 kb of the insert were sequenced starting from the following oligonucleotide primer: 5'-CGG GTC TAC ACC GCT AGC CCA GG-3'.
- 10 For identification of the insertion site lying downstream of the transposon, the chromosomal DNA of the mutant was cleaved with the restriction endonuclease XbaI and ligated in the vector pUC18 linearized with XbaI. Further cloning was carried out as described above. The nucleotide
- 15 sequence of the insertion site on one of the plasmids with an insert approx. 8.5 kb in size was determined by the dideoxy chain termination method of Sanger et al. (Proceedings of the National Academy of Sciences of the United States of America USA (1977) 74: 5463-5467). For
- 20 this, 0.65 kb of the insert were sequenced starting from the following oligonucleotide primer: 5'-CGG TGC CTT ATC CAT TCA GG-3'.

- The nucleotide sequences obtained were analysed and joined together with the Lasergene program package (Biocomputing
- 25 Software for Windows, DNASTAR, Madison, USA)). This nucleotide sequence is reproduced as SEQ ID NO 1. The result of the analysis was identification of an open reading frame of 1467 bp in length. The corresponding gene was designated the thrE gene. The associated gene product
- 30 comprises 489 amino acids and is reproduced as SEQ ID NO 2.

#### Example 2

Cloning and sequencing of the thrE gene from  
*Corynebacterium glutamicum* ATCC13032

- The thrE gene was cloned in the *E. coli* cloning vector
- 35 pUC18 (Norranders et al., Gene (1983) 26: 101-106, Roche

Diagnostics, Mannheim, Germany). The cloning was carried out in two steps. The gene from *Corynebacterium glutamicum* ATCC13032 was first amplified by a polymerase chain reaction (PCR) by means of the following oligonucleotide  
5 primers derived from SEQ ID NO 1.

thrE-forward:

5'-CCC CTT TGA CCT GGT GTT ATT G-3'

thrE-reverse:

5'-CGG CTG CGG TTT CCT CTT-3'

- 10 The PCR reaction was carried out in 30 cycles in the presence of 200  $\mu$ M deoxynucleotide triphosphates (dATP, dCTP, dGTP, dTTP), in each case 1  $\mu$ M of the corresponding oligonucleotide, 100 ng chromosomal DNA from *Corynebacterium glutamicum* ATCC13032, 1/10 volume 10-fold  
15 reaction buffer and 2.6 units of a heat-stable Taq-/Pwo-DNA polymerase mixture (Expand High Fidelity PCR System from Roche Diagnostics, Mannheim, Germany) in a Thermocycler (PTC-100, MJ Research, Inc., Watertown, USA) under the following conditions: 94°C for 30 seconds, 58°C for 30  
20 seconds and 72°C for 2 minutes.

- The amplified fragment about 1.9 kb in size was then subsequently ligated with the aid of the SureClone Ligation Kit (Amersham Pharmacia Biotech, Uppsala, Sweden) into the SmaI cleavage site of the vector pUC18 in accordance with  
25 the manufacturer's instructions. The *E. coli* strain DH5 $\alpha$ mc<sup>r</sup> (Grant et al., Proceedings of the National Academy of Sciences of the United States of America USA (1990) 87: 4645-4649) was transformed with the entire ligation batch. Transformants were identified with the aid of their  
30 carbenicillin resistance on LB agar plates containing 50  $\mu$ g/mL carbenicillin. The plasmids were prepared from 8 of the transformants and checked for the presence of the 1.9 kb PCR fragment as an insert by restriction analysis.

The recombinant plasmid formed in this way is designated pUC18thrE in the following.

The nucleotide sequence of the 1.9 kb PCR fragment in plasmid pUC18thrE was determined by the dideoxy chain  
5 termination method of Sanger et al. (Proceedings of the National Academy of Sciences of the United States of America USA (1977) 74: 5463-5467). For this, the complete insert of pUC18thrE was sequenced with the aid of the following primers from Roche Diagnostics (Mannheim,  
10 Germany).

Universal primer:

5'-GTA AAA CGA CGG CCA GT-3'

Reverse primer:

5'-GGA AAC AGC TAT GAC CAT G-3'

15 The nucleotide sequence is reproduced as SEQ ID NO 3. The nucleotide sequence obtained was analysed with the Lasergene program package (Biocomputing Software for Windows, DNASTAR, Madison, USA). The result of the analysis was identification of an open reading frame of  
20 1467 bp in length, which was designated the thrE gene. This codes for a polypeptide of 489 amino acids, which is reproduced as SEQ ID NO 4.

### Example 3

Expression of the thrE gene in *Corynebacterium glutamicum*

25 The thrE gene from *Corynebacterium glutamicum* ATCC13032 described under example 2 was cloned for expression in the vector pZ1 (Menkel et al., Applied and Environmental Microbiology (1989) 64: 549-554). For this, a DNA fragment 1881 bp in size which contained the thrE gene was cut out  
30 of the plasmid pUC18thrE with the restriction enzymes SacI and XbaI. The 5' and 3' ends of this fragment were treated with Klenow enzyme. The resulting DNA fragment was ligated

in the vector pZ1, which was linearized with ScaI and dephosphorylated beforehand. The *E. coli* strain DH5 $\alpha$ mc<sup>r</sup> (Grant et al., Proceedings of the National Academy of Sciences of the United States of America USA (1990) 87: 4645-4649) was transformed with the entire ligation batch. Transformants were identified with the aid of their kanamycin resistance on LB agar plates containing 50  $\mu$ g/mL kanamycin. The plasmids were prepared from 2 transformants and checked for the presence of the 1881 bp ScaI/XbaI fragment as an insert by restriction analysis. The recombinant plasmid formed in this manner was designated pZ1thrE (figure 2).

By means of electroporation (Haynes et al., FEMS Microbiology Letters (1989) 61: 329-334), the plasmid pZ1thrE was introduced into the threonine-forming strain *Brevibacterium flavum* DM368-2. The strain DM368-2 is described in EP-B-0 385 940 and deposited as DSM5399. Transformants were identified with the aid of their kanamycin resistance on LBHIS agar plates containing 15  $\mu$ g/mL kanamycin (Liebl et al., FEMS Microbiology Letters (1989) 65: 299-304). The strain *Brevibacterium flavum* DM368-2 pZ1thrE was formed in this manner.

#### Example 4

##### Construction of the expression plasmid pTrc99AthrE

The thrE gene from *Corynebacterium glutamicum* ATCC13032 described under example 2 was cloned for expression in *Escherichia coli* in the vector pTrc99A, which was obtained from Pharmacia Biotech (Uppsala, Sweden), for expression in *Escherichia coli*. For this, the plasmid pUC18thrE was cleaved with the enzyme SalI and the projecting 3' ends were treated with Klenow enzyme. After restriction with the enzyme KpnI, the cleavage batch was separated in 0.8% agarose gel and the thrE fragment 1.9 kbp in size was isolated with the aid of the QIAquick Gel Extraction Kit (QIAGEN, Hilden, Germany). The vector pTrc99A was cleaved with the enzyme EcoRI, the 3' ends were treated with Klenow



enzyme, cleaved with the enzyme KpnI and ligated with the thrE fragment isolated. The ligation batch was transformed in the E. coli strain DH5 $\alpha$ . Selection of cells carrying pTrc99A was carried out on LB agar (Lennox, Virology 1:190 5 (1955)), to which 50  $\mu$ g/ml ampicillin had been added. Successful cloning of the thrE gene could be demonstrated after plasmid DNA isolation and control cleavage with XbaI, BamHI, EcoRI, HindIII and SspI. The plasmid was designated pTrc99AthrE (figure 3).

#### 10 Example 5

Preparation of L-threonine with the strain MG442/pTrc99AthrE

The L-threonine-producing E. coli strain MG442 is described in the patent specification US-A- 4,278,765 and deposited 15 at the Russian National Collection of Industrial Microorganisms (VKPM, Moscow, Russia) as CMIM B-1628.

The strain MG442 was transformed with the plasmid pTrc99AthrE and plasmid-carrying cells were selected on LB agar with 50  $\mu$ g/ml ampicillin. Selected individual 20 colonies were then multiplied further on minimal medium with the following composition: 3.5 g/l Na<sub>2</sub>HPO<sub>4</sub>\*2H<sub>2</sub>O, 1.5 g/l KH<sub>2</sub>PO<sub>4</sub>, 1 g/l NH<sub>4</sub>Cl, 0.1 g/l MgSO<sub>4</sub>\*7H<sub>2</sub>O, 2 g/l glucose, 20 g/l agar, 50 mg/l ampicillin. The formation of L-threonine was checked in batch cultures of 10 ml 25 contained in 100 ml conical flasks. For this, 10 ml preculture medium of the following composition: 2 g/l yeast extract, 10 g/l (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 1 g/l KH<sub>2</sub>PO<sub>4</sub>, 0.5 g/l MgSO<sub>4</sub>\*7H<sub>2</sub>O, 15 g/l CaCO<sub>3</sub>, 20 g/l glucose, 50 mg/l ampicillin were inoculated and incubated for 16 hours at 37°C and 180 rpm 30 on an ESR incubator from Kühner AG (Birsfelden, Switzerland). 250  $\mu$ l portions of this preculture were transinoculated into 10 ml of production medium (25 g/l (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 2 g/l KH<sub>2</sub>PO<sub>4</sub>, 1 g/l MgSO<sub>4</sub>\*7H<sub>2</sub>O, 0.03 g/l FeSO<sub>4</sub>\*7H<sub>2</sub>O, 0.018 g/l MnSO<sub>4</sub>\*1H<sub>2</sub>O, 30 g/l CaCO<sub>3</sub>, 20 g/l 35 glucose) and the mixtures were incubated for 48 hours at 37°C. For induction of the expression of the thrE gene,

200 mg/l isopropyl  $\beta$ -D-thiogalactoside (IPTG) were added in parallel batches. After the incubation, the optical density (OD) of the culture suspension was determined with an LP2W photometer from Dr. Lange (Berlin, Germany) at a measurement wavelength of 660 nm.

The concentration of L-threonine formed was then determined in the sterile-filtered culture supernatant with an amino acid analyser from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column reaction with ninhydrin detection.

The result of the experiment is shown in table 2.

Table 2

Strain	Additives	OD	L-Threonine g/l
MG442	-	5.6	1.38
MG442/pTrc99AthrE	-	4.6	1.65
MG442/pTrc99AthrE	IPTG	3.6	3.5

#### Example 6

- 15 Preparation of L-threonine with the strain  
B-3996kur $\Delta$ t<sub>dh</sub>/pVIC40, pMW218thrE

The L-threonine-producing E. coli strain B-3996 is described in US-A- 5,175,107 and deposited at the Russian National Collection for Industrial Microorganisms (VKPM,  
20 Moscow, Russia).

#### 6.1 Cloning of the thrE gene in the plasmid vector pMW218

The plasmid pTrc99AthrE described under example 4 was cleaved with the enzyme SspI, the cleavage batch was separated in 0.8% agarose gel and the DNA fragment 2.5 kbp

in size, which contained the trc promoter region and the rRNA terminator region in addition to the thrE gene, was isolated with the aid of the "QIAquick Gel Extraction Kit" (QIAGEN, Hilden, Germany). The plasmid pMW218 (Nippon Gene, Toyama, Japan) was cleaved with the enzyme SmaI and ligated with the thrE fragment. The E. coli strain DH5 $\alpha$  was transformed with the ligation batch and pMW218-carrying cells were selected by plating out on LB agar, to which 20  $\mu$ g/ml kanamycin are added. Successful cloning of the thrE gene could be demonstrated after plasmid DNA isolation and control cleavage with HindIII and ClaI. The plasmid was designated pMW218thrE (figure 4).

#### 6.2 Preparation of the strain B-3996kur $\Delta$ tdh/pVIC40, pMW218thrE

- After culture in antibiotic-free complete medium for approximately ten generations, a derivative of strain B-3996 which no longer contained the plasmid pVIC40 was isolated. The strain formed was streptomycin-sensitive and was designated B-3996kur.
- The method described by Hamilton et al. (Journal of Bacteriology (1989) 171: 4617-4622), which is based on the use of the plasmid pMAK705 with a temperature-sensitive replicon, was used for incorporation of a deletion into the tdh gene. The plasmid pDR121 (Ravnikar and Somerville, Journal of Bacteriology (1987) 169: 4716-4721) contains a DNA fragment from E. coli 3.7 kilo-base pairs (kbp) in size, on which the tdh gene is coded. To generate a deletion of the tdh gene region, pDR121 was cleaved with the restriction enzymes ClaI and EcoRV and the DNA fragment 5 kbp in size isolated was ligated, after treatment with Klenow enzyme. The ligation batch was transformed in the E. coli strain DH5 $\alpha$  and plasmid-carrying cells were selected on LB agar, to which 50  $\mu$ g/ml ampicillin are added.
- Successful deletion of the tdh gene could be demonstrated after plasmid DNA isolation and control cleavage with

EcoRI. The EcoRI fragment 1.7 kbp in size was isolated, and ligated with the plasmid pMAK705, which was partly digested with EcoRI. The ligation batch was transformed in DH5 $\alpha$  and plasmid-carrying cells were selected on LB agar, 5 to which 20  $\mu$ g/ml chloramphenicol were added. Successful cloning was demonstrated after isolation of the plasmid DNA and cleavage with EcoRI. The pMAK705 derivative formed was designated pDM32.

For the gene replacement, B-3996kur was transformed with 10 the plasmid pDM32. The replacement of the chromosomal tdh gene with the plasmid-coded deletion construct was carried out by the selection process described by Hamilton et al. and was verified by standard PCR methods (Innis et al. (1990), PCR Protocols. A Guide to Methods and Applications, 15 Academic Press) with the following oligonucleotide primers:

Tdh1: 5'-TCGCGACCTATAAGTTTGGG-3'

Tdh2: 5'-AATACCAGCCCTTGTTTCGTG-3'

The strain formed was tested for kanamycin sensitivity and was designated B-3996kur $\Delta$ tdh.

20 B-3996kur $\Delta$ tdh was transformed with the plasmid pVIC40 isolated from B-3996 and plasmid-carrying cells were selected on LB agar supplemented with 20  $\mu$ g/ml streptomycin. A selected individual colony was designated B-3996kur $\Delta$ tdh/pVIC40 and transformed with the plasmid 25 pMW218thrE. Selection is carried out on LB-agar to which 20  $\mu$ g/ml streptomycin and 50  $\mu$ g/ml kanamycin are added. The strain formed in this way was designated B-3996kur $\Delta$ tdh/pVIC40, pMW218thrE.

### 6.3 Preparation of L-threonine

30 The preparation of L-threonine by the strains B-3996kur $\Delta$ tdh/pVIC40 and B-3996kur $\Delta$ tdh/pVIC40, pMW218thrE was tested as described in example 5. The minimal medium, the preculture medium and the production medium were supplemented with 20  $\mu$ g/ml streptomycin for

B-3996kur $\Delta$ tdh/pVIC40 and with 20  $\mu$ g/ml streptomycin and 50  $\mu$ g/ml kanamycin for B-3996kur $\Delta$ tdh/pVIC40, pMW218thrE.

The result of the experiment is summarized in table 3.

Table 3

Strain	OD (660 nm)	L-Threonine g/l
B-3996kur $\Delta$ tdh/pVIC40	4.7	6.26
B-3996kur $\Delta$ tdh/pVIC40, pMW218thrE	4.8	7.57

5

The following figures are attached:

- Figure 1: Map of the plasmid pCGL0040 containing the transposon Tn5531. The transposon is identified as the non-shaded arrow.
- 5 • Figure 2: Map of the plasmid pZ1thrE containing the thrE gene.
- Figure 3: Map of the plasmid pTrc99AthrE containing the thrE gene.
- Figure 4: Map of the plasmid pMW218thrE containing the  
10 thrE gene

The length data are to be understood as approx. data. The abbreviations and designations used have the following meaning:

- Amp: Ampicillin resistance gene
- 15 • Kan: Kanamycin resistance gene
- 'amp: 3' part of the ampicillin resistance gene
- oriBR322: Replication region of the plasmid pBR322
- lacI: Gene for the repressor protein of the trc promoter
- 20 • P<sub>trc</sub>: trc promoter region, IPTG-inducible
- 5S: 5S rRNA region
- rrnBT: rRNA terminator region

The abbreviations for the restriction enzymes have the following meaning.

- 25 • BamHI: Restriction endonuclease from *Bacillus amyloliquefaciens*
- BglII: Restriction endonuclease from *Bacillus globigii*


- ClaI: Restriction endonuclease from *Caryophanon latum*
- EcoRI: Restriction endonuclease from *Escherichia coli*
- EcoRV: Restriction endonuclease from *Escherichia coli*
- HindIII: Restriction endonuclease from *Haemophilus*  
5     *influenzae*
- KpnI: Restriction endonuclease from *Klebsiella*  
          *pneumoniae*
- PstI: Restriction endonuclease from *Providencia stuartii*
- PvuI: Restriction endonuclease from *Proteus vulgaris*
- 10 • SacI: Restriction endonuclease from *Streptomyces*  
          *achromogenes*
- SalI: Restriction endonuclease from *Streptomyces albus*
- SmaI: Restriction endonuclease from *Serratia marcescens*
- XbaI: Restriction endonuclease from *Xanthomonas badrii*
- 15 • XhoI: Restriction endonuclease from *Xanthomonas*  
          *holcicola*

Original (for SUBMISSION) - printed on 04.04.2001 02:03:45 PM

0-1	Form - PCT/RO/134 (EASY) Indications Relating to Deposited Microorganism(s) or Other Biological Material (PCT Rule 13bis)	
0-1-1	Prepared using	PCT-EASY Version 2.91 (updated 01.01.2001)
0-2	International Application No.	
0-3	Applicant's or agent's file reference	000225 BT

1	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
1-1	page	8
1-2	line	1-7
1-3	Identification of Deposit	
1-3-1	Name of depositary institution	DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH
1-3-2	Address of depositary institution	Mascheroder Weg 1b, D-38124 Braunschweig, Germany
1-3-3	Date of deposit	03 June 1999 (03.06.1999)
1-3-4	Accession Number	DSMZ 12840
1-4	Additional Indications	NONE
1-5	Designated States for Which Indications are Made	all designated States
1-6	Separate Furnishing of Indications  These indications will be submitted to the International Bureau later	NONE

## FOR RECEIVING OFFICE USE ONLY

0-4	This form was received with the international application: (yes or no)	YES
0-4-1	Authorized officer	 C.A.J.A. FIEBIGER

## FOR INTERNATIONAL BUREAU USE ONLY

0-5	This form was received by the international Bureau on:	
0-5-1	Authorized officer	



## Patent claims

1. A process for the fermentative preparation of L-threonine, which comprises employing  
5 Enterobacteriaceae bacteria, in particular those which already produce L-threonine and in which the nucleotide sequence(s) of coryneform bacteria which code(s) for the thrE gene are enhanced, in particular over-expressed.
- 10 2. A process as claimed in claim 1, wherein further genes are enhanced in addition to the thrE gene.
3. A process as claimed in claim 1 or 2, wherein the microorganisms of the family Enterobacteriaceae are from the genus Escherichia and Serratia.
- 15 4. A process as claimed in claim 3, wherein the microorganisms are from the genus Escherichia , in particular of the species Escherichia coli.
5. A process as claimed in claim 1, wherein the thrABC operon which codes for aspartate kinase, homoserine  
20 dehydrogenase, homoserine kinase and threonine synthase is enhanced at the same time.
6. A process as claimed in claim 1, wherein the pyc gene which codes for pyruvate carboxylase is enhanced at the same time.
- 25 7. A process as claimed in claim 1, wherein the pps gene which codes for phosphoenol pyruvate synthase is enhanced at the same time.
8. A process as claimed in claim 1, wherein the ppc gene which codes for phosphoenol pyruvate carboxylase is  
30 enhanced at the same time.

9. A process as claimed in claim 1, wherein the genes pntA and pntB which code for transhydrogenase are enhanced at the same time.
10. A process as claimed in claim 1, wherein bacteria in which the metabolic pathways which reduce the formation of L-threonine are at least partly eliminated are employed.
11. A process as claimed in claim 1, wherein a strain transformed with a plasmid vector is employed and the plasmid vector carries the nucleotide sequence which codes for the thrE gene of coryneform bacteria.
12. A process as claimed in claim 1, wherein bacteria transformed with the plasmid pZ1thrE are employed.
13. A process as claimed in claim 1, wherein the expression of the thrE gene is induced, in particular with isopropyl  $\beta$ -D-thiogalactoside.
14. A process as claimed in claim 1, wherein the gdhA gene which codes for glutamate dehydrogenase is enhanced at the same time.
15. A process as claimed in claim 1, wherein the rhtB gene which imparts homoserine resistance is enhanced at the same time.
16. A process for the preparation of L-threonine, which comprises carrying out the following steps:
  - a) fermentation of microorganisms of the family Enterobacteriaceae in which at least the thrE gene of coryneform bacteria is enhanced (over-expressed), optionally in combination with further genes,
  - b) concentration of the L-threonine in the medium or in the cells of the microorganisms of the family Enterobacteriaceae, and

c) isolation of the L-threonine.

17. The plasmid pZ1thrE which contains the thrE gene of *Corynebacterium glutamicum* ATCC13032.
18. The *Brevibacterium flavum* strain DM368-2 pZ1thrE  
5 deposited as DSM 12840 at the DSMZ [German Collection  
of Microorganisms and Cell Cultures], Braunschweig.

Figure 1




Figure 2




Figure 3





Figure 4



## SEQUENCE PROTOCOL

&lt;110&gt; Degussa AG

5 &lt;120&gt; Process for the fermentative preparation of L-threonine.

&lt;130&gt; 000225 BT

&lt;140&gt;

10 &lt;141&gt;

&lt;160&gt; 4

&lt;170&gt; PatentIn Ver. 2.1

15

&lt;210&gt; 1

&lt;211&gt; 2817

&lt;212&gt; DNA

&lt;213&gt; Corynebacterium glutamicum ATCC14752

20

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (398)..(1864)

&lt;223&gt; thrE gene

25

&lt;400&gt; 1

aatgaaataa tcccctcacc aactggcgac attcaaacac cgtttcattt ccaaacatcg 60

agccaaggga aaagaaagcc cctaagcccc gtgttattaa atggagactc tttggagacc 120

30

tcaagccaaa aaggggcatt ttcattaaga aaatacccct ttgacctggg gttattgagc 180

tggaagaagag acttgaactc tcaacctacg cattacaagt gcgttgcgct gccaatgtcg 240

35

ccactccagc accgcagatg ctgatgatca acaactacga atacgtatct tagcgtatgt 300

gtacatcaca atggaattcg gggctagagt atctgggtgaa ccgtgcataa acgacctgtg 360

attggactct ttttccttgc aaaatgtttt ccagcgg atg ttg agt ttt gcg acc 415

40

Met Leu Ser Phe Ala Thr  
1 5

ctt cgt ggc cgc att tca aca gtt gac gct gca aaa gcc gca cct ccg 463

Leu Arg Gly Arg Ile Ser Thr Val Asp Ala Ala Lys Ala Ala Pro Pro  
10 15 20

45

cca tcg cca cta gcc ccg att gat ctc act gac cat agt caa gtg gcc 511

Pro Ser Pro Leu Ala Pro Ile Asp Leu Thr Asp His Ser Gln Val Ala  
25 30 35

50

ggg gtg atg aat ttg gct gcg aga att ggc gat att ttg ctt tct tca 559

Gly Val Met Asn Leu Ala Ala Arg Ile Gly Asp Ile Leu Leu Ser Ser  
40 45 50

ggg acg tca aac agt gat acc aag gtg caa gtt cga gcg gtg acc tct 607

Gly Thr Ser Asn Ser Asp Thr Lys Val Gln Val Arg Ala Val Thr Ser  
55 60 65 70

gcg tat ggc ctg tac tat acg cat gtg gat atc acg ttg aat acg atc 655

Ala Tyr Gly Leu Tyr Tyr Thr His Val Asp Ile Thr Leu Asn Thr Ile  
75 80 85

60

acc atc ttc acc aac atc ggt gtg gag agg aag atg ccg gtc aac gtg 703

Thr Ile Phe Thr Asn Ile Gly Val Glu Arg Lys Met Pro Val Asn Val  
90 95 100

65



	ttt	cat	gtt	gtg	ggc	aag	ttg	gac	acc	aac	ttc	tcc	aaa	ctg	tct	gag	751
	Phe	His	Val	Val	Gly	Lys	Leu	Asp	Thr	Asn	Phe	Ser	Lys	Leu	Ser	Glu	
			105					110					115				
5	gtt	gac	cgt	ttg	atc	cgt	tcc	att	cag	gct	ggt	gct	acc	ccg	cct	gag	799
	Val	Asp	Arg	Leu	Ile	Arg	Ser	Ile	Gln	Ala	Gly	Ala	Thr	Pro	Pro	Glu	
		120					125					130					
10	gtt	gcc	gag	aaa	att	ctg	gac	gag	ttg	gag	caa	tcg	cct	gcg	tct	tat	847
	Val	Ala	Glu	Lys	Ile	Leu	Asp	Glu	Leu	Glu	Gln	Ser	Pro	Ala	Ser	Tyr	
	135					140					145					150	
15	ggt	ttc	cct	gtt	gcg	ttg	ctt	ggc	tgg	gca	atg	atg	ggt	ggc	gct	gtt	895
	Gly	Phe	Pro	Val	Ala	Leu	Leu	Gly	Trp	Ala	Met	Met	Gly	Gly	Ala	Val	
					155					160					165		
20	gct	gtg	ctg	ttg	ggt	ggt	gga	tgg	cag	gtt	tcc	cta	att	gct	ttt	att	943
	Ala	Val	Leu	Leu	Gly	Gly	Gly	Trp	Gln	Val	Ser	Leu	Ile	Ala	Phe	Ile	
				170					175					180			
	acc	gcg	ttc	acg	atc	att	gcc	acg	acg	tca	ttt	ttg	gga	aag	aag	ggt	991
	Thr	Ala	Phe	Thr	Ile	Ile	Ala	Thr	Thr	Ser	Phe	Leu	Gly	Lys	Lys	Gly	
			185					190					195				
25	ttg	cct	act	ttc	ttc	caa	aat	gtt	gtt	ggt	ggt	ttt	att	gcc	acg	ctg	1039
	Leu	Pro	Thr	Phe	Phe	Gln	Asn	Val	Val	Gly	Gly	Phe	Ile	Ala	Thr	Leu	
		200					205					210					
30	cct	gca	tcg	att	gct	tat	tct	ttg	gcg	ttg	caa	ttt	ggt	ctt	gag	atc	1087
	Pro	Ala	Ser	Ile	Ala	Tyr	Ser	Leu	Ala	Leu	Gln	Phe	Gly	Leu	Glu	Ile	
	215					220					225					230	
35	aaa	ccg	agc	cag	atc	atc	gca	tct	gga	att	gtt	gtg	ctg	ttg	gca	ggt	1135
	Lys	Pro	Ser	Gln	Ile	Ile	Ala	Ser	Gly	Ile	Val	Val	Leu	Leu	Ala	Gly	
				235						240					245		
40	ttg	aca	ctt	gtg	caa	tct	ctg	cag	gac	ggc	atc	acg	ggc	gct	ccg	gtg	1183
	Leu	Thr	Leu	Val	Gln	Ser	Leu	Gln	Asp	Gly	Ile	Thr	Gly	Ala	Pro	Val	
				250					255					260			
	aca	gca	agt	gca	cga	ttt	ttt	gaa	aca	ctc	ctg	ttt	acc	ggc	ggc	att	1231
	Thr	Ala	Ser	Ala	Arg	Phe	Phe	Glu	Thr	Leu	Leu	Phe	Thr	Gly	Gly	Ile	
			265					270					275				
45	gtt	gct	ggc	gtg	ggt	ttg	ggc	att	cag	ctt	tct	gaa	atc	ttg	cat	gtc	1279
	Val	Ala	Gly	Val	Gly	Leu	Gly	Ile	Gln	Leu	Ser	Glu	Ile	Leu	His	Val	
		280				285						290					
50	atg	ttg	cct	gcc	atg	gag	tcc	gct	gca	gca	cct	aat	tat	tcg	tct	aca	1327
	Met	Leu	Pro	Ala	Met	Glu	Ser	Ala	Ala	Ala	Pro	Asn	Tyr	Ser	Ser	Thr	
	295					300					305					310	
55	ttc	gcc	cgc	att	atc	gct	ggt	ggc	gtc	acc	gca	gcg	gcc	ttc	gca	gtg	1375
	Phe	Ala	Arg	Ile	Ile	Ala	Gly	Gly	Val	Thr	Ala	Ala	Ala	Phe	Ala	Val	
				315						320					325		
	ggt	tgt	tac	gcg	gag	tgg	tcc	tcg	gtg	att	att	gcg	ggg	ctt	act	gcg	1423
	Gly	Cys	Tyr	Ala	Glu	Trp	Ser	Ser	Val	Ile	Ile	Ala	Gly	Leu	Thr	Ala	
				330					335					340			

ctg atg ggt tct gcg ttt tat tac ctc ttc gtt gtt tat tta ggc ccc 1471  
 Leu Met Gly Ser Ala Phe Tyr Tyr Leu Phe Val Val Tyr Leu Gly Pro  
 345 350 355

5 gtc tct gcc gct gcg att gct gca aca gca gtt ggt ttc act ggt ggt 1519  
 Val Ser Ala Ala Ala Ile Ala Ala Thr Ala Val Gly Phe Thr Gly Gly  
 360 365 370

10 ttg ctt gcc cgt cga ttc ttg att cca ccg ttg att gtg gcg att gcc 1567  
 Leu Leu Ala Arg Arg Phe Leu Ile Pro Pro Leu Ile Val Ala Ile Ala  
 375 380 385 390

15 ggc atc aca cca atg ctt cca ggt cta gca att tac cgc gga atg tac 1615  
 Gly Ile Thr Pro Met Leu Pro Gly Leu Ala Ile Tyr Arg Gly Met Tyr  
 395 400 405

20 gcc acc ttg aat gat caa aca ctc atg ggt ttc acc aac att gcg gtt 1663  
 Ala Thr Leu Asn Asp Gln Thr Leu Met Gly Phe Thr Asn Ile Ala Val  
 410 415 420

25 gct tta gcc act gct tca tca ctt gcc gct gcc gtg gtt ttg ggt gag 1711  
 Ala Leu Ala Thr Ala Ser Ser Leu Ala Ala Gly Val Val Leu Gly Glu  
 425 430 435

tgg att gcc cgt agg cta cgt cgt cca cca cgc ttc aac cca tac cgt 1759  
 Trp Ile Ala Arg Arg Leu Arg Arg Pro Pro Arg Phe Asn Pro Tyr Arg  
 440 445 450

30 gca ttt acc aag gcg aat gag ttc tcc ttc cag gag gaa gct gag cag 1807  
 Ala Phe Thr Lys Ala Asn Glu Phe Ser Phe Glu Glu Ala Glu Gln  
 455 460 465 470

35 aat cag cgc cgg cag aga aaa cgt cca aag act aat caa aga ttc ggt 1855  
 Asn Gln Arg Arg Gln Arg Lys Arg Pro Lys Thr Asn Gln Arg Phe Gly  
 475 480 485

40 aat aaa agg taaaaatcaa cctgcttagg cgtctttcgc ttaaatagcg 1904  
 Asn Lys Arg

tagaatatcg ggtcgatcgc ttttaaacac tcaggaggat ccttgccggc caaaatcacg 1964  
 gacactcgtc ccaccccaga atcccttcac gctgttgaag aggaaaccgc agccggtgcc 2024

45 cgcaggattg ttgccaccta ttctaaggac ttcttcgacg gcgtcacttt gatgtgcatg 2084  
 ctccggcgttg aacctcaggg cctgcgttac accaaggctc cttctgaaca cgaggaagct 2144  
 cagccaaaga aggctacaaa gcggactcgt aaggcaccag ctaagaaggc tgctgctaag 2204

50 aaaacgacca agaagaccac taagaaaact actaaaaaga ccaccgcaaa gaagaccaca 2264  
 aagaagtctt aagccggatc ttatatggat gattccaata gctttgtagt tgttgctaac 2324

55 cgtctgccag tggatatgac tgtccaccca gatggttagct atagcatctc ccccgcccc 2384  
 ggtggccttg tcacggggct tcccccggt ctggaacaac atcgtggatg ttgggtcgga 2444  
 tggcctggaa ctgtagatgt tgcaccogaa ccatttcgaa cagatacggg tgttttgctg 2504

60 caccctgttg tcctcactgc aagtgactat gaaggcttct acgagggctt ttcaaacgca 2564  
 acgctgtggc ctcttttcca cgatttgatt gttactccgg tgtacaacac cgattggtgg 2624

65 catgcgtttc gggaagtaaa cctcaagttc gctgaagccg tgagccaagt ggcggcacac 2684

ggtgccactg tgtgggtgca ggactatcag ctgttgctgg ttcttgcat tttgcgccag 2744  
 atgcgccctg atttgaagat cggtttcttc ctccacattc ccttcccttc cctgatctg 2804  
 5 ttccgtcagc tgc 2817

<210> 2  
 <211> 489  
 10 <212> PRT  
 <213> Corynebacterium glutamicum ATCC14752

<400> 2  
 15 Met Leu Ser Phe Ala Thr Leu Arg Gly Arg Ile Ser Thr Val Asp Ala  
 1 5 10 15  
 Ala Lys Ala Ala Pro Pro Pro Ser Pro Leu Ala Pro Ile Asp Leu Thr  
 20 20 25 30  
 20 Asp His Ser Gln Val Ala Gly Val Met Asn Leu Ala Ala Arg Ile Gly  
 35 40 45  
 Asp Ile Leu Leu Ser Ser Gly Thr Ser Asn Ser Asp Thr Lys Val Gln  
 50 55 60  
 25 Val Arg Ala Val Thr Ser Ala Tyr Gly Leu Tyr Tyr Thr His Val Asp  
 65 70 75 80  
 Ile Thr Leu Asn Thr Ile Thr Ile Phe Thr Asn Ile Gly Val Glu Arg  
 85 90 95  
 Lys Met Pro Val Asn Val Phe His Val Val Gly Lys Leu Asp Thr Asn  
 100 105 110  
 35 Phe Ser Lys Leu Ser Glu Val Asp Arg Leu Ile Arg Ser Ile Gln Ala  
 115 120 125  
 Gly Ala Thr Pro Pro Glu Val Ala Glu Lys Ile Leu Asp Glu Leu Glu  
 130 135 140  
 40 Gln Ser Pro Ala Ser Tyr Gly Phe Pro Val Ala Leu Leu Gly Trp Ala  
 145 150 155 160  
 Met Met Gly Gly Ala Val Ala Val Leu Leu Gly Gly Gly Trp Gln Val  
 165 170 175  
 Ser Leu Ile Ala Phe Ile Thr Ala Phe Thr Ile Ile Ala Thr Thr Ser  
 180 185 190  
 50 Phe Leu Gly Lys Lys Gly Leu Pro Thr Phe Phe Gln Asn Val Val Gly  
 195 200 205  
 Gly Phe Ile Ala Thr Leu Pro Ala Ser Ile Ala Tyr Ser Leu Ala Leu  
 210 215 220  
 55 Gln Phe Gly Leu Glu Ile Lys Pro Ser Gln Ile Ile Ala Ser Gly Ile  
 225 230 235 240  
 Val Val Leu Leu Ala Gly Leu Thr Leu Val Gln Ser Leu Gln Asp Gly  
 245 250 255  
 60 Ile Thr Gly Ala Pro Val Thr Ala Ser Ala Arg Phe Phe Glu Thr Leu  
 260 265 270  
 65 Leu Phe Thr Gly Gly Ile Val Ala Gly Val Gly Leu Gly Ile Gln Leu  
 275 280 285

Ser Glu Ile Leu His Val Met Leu Pro Ala Met Glu Ser Ala Ala Ala  
 290 295 300  
 5 Pro Asn Tyr Ser Ser Thr Phe Ala Arg Ile Ile Ala Gly Gly Val Thr  
 305 310 315 320  
 Ala Ala Ala Phe Ala Val Gly Cys Tyr Ala Glu Trp Ser Ser Val Ile  
 325 330 335  
 10 Ile Ala Gly Leu Thr Ala Leu Met Gly Ser Ala Phe Tyr Tyr Leu Phe  
 340 345 350  
 Val Val Tyr Leu Gly Pro Val Ser Ala Ala Ala Ile Ala Ala Thr Ala  
 355 360 365  
 15 Val Gly Phe Thr Gly Gly Leu Leu Ala Arg Arg Phe Leu Ile Pro Pro  
 370 375 380  
 20 Leu Ile Val Ala Ile Ala Gly Ile Thr Pro Met Leu Pro Gly Leu Ala  
 385 390 395 400  
 Ile Tyr Arg Gly Met Tyr Ala Thr Leu Asn Asp Gln Thr Leu Met Gly  
 405 410 415  
 25 Phe Thr Asn Ile Ala Val Ala Leu Ala Thr Ala Ser Ser Leu Ala Ala  
 420 425 430  
 Gly Val Val Leu Gly Glu Trp Ile Ala Arg Arg Leu Arg Arg Pro Pro  
 435 440 445  
 30 Arg Phe Asn Pro Tyr Arg Ala Phe Thr Lys Ala Asn Glu Phe Ser Phe  
 450 455 460  
 35 Gln Glu Glu Ala Glu Gln Asn Gln Arg Arg Gln Arg Lys Arg Pro Lys  
 465 470 475 480  
 Thr Asn Gln Arg Phe Gly Asn Lys Arg  
 485  
 40  
 <210> 3  
 <211> 1909  
 45 <212> DNA  
 <213> Corynebacterium glutamicum ATCC13032  
 <220>  
 <221> CDS  
 50 <222> (280)..(1746)  
 <223> thrE gene  
 <400> 3  
 agcttgcatg cctgcaggtc gactctagag gatccccccc ctttgacctg gtgttattga 60  
 55 gctggagaag agacttgaac tctcaacctt cgcattacaa gtgcgttgcg ctgccaatg 120  
 cgccactcca gcaccgcaga tgctgatgat caacaactac gaatacgtat cttagcgtat 180  
 60 gtgtacatca caatggaatt cggggctaga gtatctggtg aaccgtgcat aaacgacctg 240

	tgattggact ctttttcctt gcaaaatgtt ttccagcgg atg ttg agt ttt gcg	294
	Met Leu Ser Phe Ala	
5	1 5	
	acc ctt cgt ggc cgc att tca aca gtt gac gct gca aaa gcc gca cct	342
	Thr Leu Arg Gly Arg Ile Ser Thr Val Asp Ala Ala Lys Ala Ala Pro	
	10 15 20	
10	ccg cca tcg cca cta gcc ccg att gat ctc act gac cat agt caa gtg	390
	Pro Pro Ser Pro Leu Ala Pro Ile Asp Leu Thr Asp His Ser Gln Val	
	25 30 35	
15	gcc ggt gtg atg aat ttg gct gcg aga att ggc gat att ttg ctt tct	438
	Ala Gly Val Met Asn Leu Ala Arg Ile Gly Asp Ile Leu Leu Ser	
	40 45 50	
20	tca ggt acg tca aat agt gac acc aag gta caa gtt cga gca gtg acc	486
	Ser Gly Thr Ser Asn Ser Asp Thr Lys Val Gln Val Arg Ala Val Thr	
	55 60 65	
	tct gcg tac ggt ttg tac tac acg cac gtg gat atc acg ttg aat acg	534
	Ser Ala Tyr Gly Leu Tyr Tyr Thr His Val Asp Ile Thr Leu Asn Thr	
	70 75 80 85	
25	atc acc atc ttc acc aac atc ggt gtg gag agg aag atg ccg gtc aac	582
	Ile Thr Ile Phe Thr Asn Ile Gly Val Glu Arg Lys Met Pro Val Asn	
	90 95 100	
30	gtg ttt cat gtt gta gcc aag ttg gac acc aac ttc tcc aaa ctg tct	630
	Val Phe His Val Gly Lys Leu Asp Thr Asn Phe Ser Lys Leu Ser	
	105 110 115	
35	gag gtt gac cgt ttg atc cgt tcc att cag gct ggt gcg acc ccg cct	678
	Glu Val Asp Arg Leu Ile Arg Ser Ile Gln Ala Gly Ala Thr Pro Pro	
	120 125 130	
40	gag gtt gcc gag aaa atc ctg gac gag ttg gag caa tcc cct gcg tct	726
	Glu Val Ala Glu Lys Ile Leu Asp Glu Leu Glu Gln Ser Pro Ala Ser	
	135 140 145	
	tat ggt ttc cct gtt gcg ttg ctt ggc tgg gca atg atg ggt ggt gct	774
	Tyr Gly Phe Pro Val Ala Leu Leu Gly Trp Ala Met Met Gly Gly Ala	
	150 155 160 165	
45	gtt gct gtg ctg ttg ggt ggt gga tgg cag gtt tcc cta att gct ttt	822
	Val Ala Val Leu Leu Gly Gly Gly Trp Gln Val Ser Leu Ile Ala Phe	
	170 175 180	
50	att acc gcg ttc acg atc att gcc acg acg tca ttt ttg gga aag aag	870
	Ile Thr Ala Phe Thr Ile Ile Ala Thr Thr Ser Phe Leu Gly Lys Lys	
	185 190 195	
55	ggt ttg cct act ttc ttc caa aat gtt gtt ggt ggt ttt att gcc acg	918
	Gly Leu Pro Thr Phe Phe Gln Asn Val Val Gly Gly Phe Ile Ala Thr	
	200 205 210	
	ctg cct gca tcg att gct tat tct ttg gcg ttg caa ttt ggt ctt gag	966
	Leu Pro Ala Ser Ile Ala Tyr Ser Leu Ala Leu Gln Phe Gly Leu Glu	
	215 220 225	
60	atc aaa ccg agc cag atc atc gca tct gga att gtt gtg ctg ttg gca	1014
	Ile Lys Pro Ser Gln Ile Ile Ala Ser Gly Ile Val Val Leu Leu Ala	
	230 235 240 245	
65		

	ggt ttg aca ctc gtg caa tct ctg cag gac ggc atc acg ggc gct ccg	1062
	Gly Leu Thr Leu Val Gln Ser Leu Gln Asp Gly Ile Thr Gly Ala Pro	
	250 255 260	
5	gtg aca gca agt gca cga ttt ttc gaa aca ctc ctg ttt acc ggc ggc	1110
	Val Thr Ala Ser Ala Arg Phe Phe Glu Thr Leu Leu Phe Thr Gly Gly	
	265 270 275	
10	att gtt gct ggc gtg ggt ttg ggc att cag ctt tct gaa atc ttg cat	1158
	Ile Val Ala Gly Val Gly Leu Gly Ile Gln Leu Ser Glu Ile Leu His	
	280 285 290	
15	gtc atg ttg cct gcc atg gag tcc gct gca gca cct aat tat tcg tct	1206
	Val Met Leu Pro Ala Met Glu Ser Ala Ala Pro Asn Tyr Ser Ser	
	295 300 305	
20	aca ttc gcc cgc att atc gct ggt ggc gtc acc gca gcg gcc ttc gca	1254
	Thr Phe Ala Arg Ile Ile Ala Gly Gly Val Thr Ala Ala Ala Phe Ala	
	310 315 320 325	
	gtg ggt tgt tac gcg gag tgg tcc tcg gtg att att gcg ggg ctt act	1302
	Val Gly Cys Tyr Ala Glu Trp Ser Ser Val Ile Ile Ala Gly Leu Thr	
	330 335 340	
25	gcg ctg atg ggt tct gcg ttt tat tac ctc ttc gtt gtt tat tta ggc	1350
	Ala Leu Met Gly Ser Ala Phe Tyr Tyr Leu Phe Val Val Tyr Leu Gly	
	345 350 355	
30	ccc gtc tct gcc gct gcg att gct gca aca gca gtt ggt ttc act ggt	1398
	Pro Val Ser Ala Ala Ala Ile Ala Ala Thr Ala Val Gly Phe Thr Gly	
	360 365 370	
35	ggt ttg ctt gcc cgt cga ttc ttg att cca ccg ttg att gtg gcg att	1446
	Gly Leu Leu Ala Arg Arg Phe Leu Ile Pro Pro Leu Ile Val Ala Ile	
	375 380 385	
40	gcc ggc atc aca cca atg ctt cca ggt cta gca att tac cgc gga atg	1494
	Ala Gly Ile Thr Pro Met Leu Pro Gly Leu Ala Ile Tyr Arg Gly Met	
	390 395 400 405	
	tac gcc acc ctg aat gat caa aca ctc atg ggt ttc acc aac att gcg	1542
	Tyr Ala Thr Leu Asn Asp Gln Thr Leu Met Gly Phe Thr Asn Ile Ala	
	410 415 420	
45	gtt gct tta gcc act gct tca tca ctt gcc gct ggc gtg gtt ttg ggt	1590
	Val Ala Leu Ala Thr Ala Ser Ser Leu Ala Ala Gly Val Val Leu Gly	
	425 430 435	
50	gag tgg att gcc cgc agg cta cgt cgt cca cca cgc ttc aac cca tac	1638
	Glu Trp Ile Ala Arg Arg Leu Arg Arg Pro Pro Arg Phe Asn Pro Tyr	
	440 445 450	
55	cgt gca ttt acc aag gcg aat gag ttc tcc ttc cag gag gaa gct gag	1686
	Arg Ala Phe Thr Lys Ala Asn Glu Phe Ser Phe Gln Glu Glu Ala Glu	
	455 460 465	
60	cag aat cag cgc cgg cag aga aaa cgt cca aag act aat cag aga ttc	1734
	Gln Asn Gln Arg Arg Gln Arg Lys Arg Pro Lys Thr Asn Gln Arg Phe	
	470 475 480 485	
	ggt aat aaa agg taaaaatcaa cctgcttagg cgtctttcgc ttaaatagcg	1786
	Gly Asn Lys Arg	
65	tagaatatcg ggtcgatcgc ttttaaacac tcaggaggat ccttgccggc caaatcacg	1846

gacactcgtc ccaccccaga atcccttcac gctgttgaag aggaaaccgc agccggggta 1906  
 ccg 1909

5  
 <210> 4  
 <211> 489  
 <212> PRT  
 <213> Corynebacterium glutamicum ATCC13032

10  
 <400> 4  
 Met Leu Ser Phe Ala Thr Leu Arg Gly Arg Ile Ser Thr Val Asp Ala  
 1 5 10 15

15 Ala Lys Ala Ala Pro Pro Pro Ser Pro Leu Ala Pro Ile Asp Leu Thr  
 20 25 30

Asp His Ser Gln Val Ala Gly Val Met Asn Leu Ala Ala Arg Ile Gly  
 35 40 45

20 Asp Ile Leu Leu Ser Ser Gly Thr Ser Asn Ser Asp Thr Lys Val Gln  
 50 55 60

Val Arg Ala Val Thr Ser Ala Tyr Gly Leu Tyr Tyr Thr His Val Asp  
 25 65 70 75 80

Ile Thr Leu Asn Thr Ile Thr Ile Phe Thr Asn Ile Gly Val Glu Arg  
 85 90 95

30 Lys Met Pro Val Asn Val Phe His Val Val Gly Lys Leu Asp Thr Asn  
 100 105 110

Phe Ser Lys Leu Ser Glu Val Asp Arg Leu Ile Arg Ser Ile Gln Ala  
 115 120 125

35 Gly Ala Thr Pro Pro Glu Val Ala Glu Lys Ile Leu Asp Glu Leu Glu  
 130 135 140

Gln Ser Pro Ala Ser Tyr Gly Phe Pro Val Ala Leu Leu Gly Trp Ala  
 40 145 150 155 160

Met Met Gly Gly Ala Val Ala Val Leu Leu Gly Gly Gly Trp Gln Val  
 165 170 175

45 Ser Leu Ile Ala Phe Ile Thr Ala Phe Thr Ile Ile Ala Thr Thr Ser  
 180 185 190

Phe Leu Gly Lys Lys Gly Leu Pro Thr Phe Phe Gln Asn Val Val Gly  
 195 200 205

50 Gly Phe Ile Ala Thr Leu Pro Ala Ser Ile Ala Tyr Ser Leu Ala Leu  
 210 215 220

Gln Phe Gly Leu Glu Ile Lys Pro Ser Gln Ile Ile Ala Ser Gly Ile  
 55 225 230 235 240

Val Val Leu Leu Ala Gly Leu Thr Leu Val Gln Ser Leu Gln Asp Gly  
 245 250 255

60 Ile Thr Gly Ala Pro Val Thr Ala Ser Ala Arg Phe Phe Glu Thr Leu  
 260 265 270

Leu Phe Thr Gly Gly Ile Val Ala Gly Val Gly Leu Gly Ile Gln Leu  
 275 280 285

65 Ser Glu Ile Leu His Val Met Leu Pro Ala Met Glu Ser Ala Ala Ala

	290		295		300												
	Pro	Asn	Tyr	Ser	Ser	Thr	Phe	Ala	Arg	Ile	Ile	Ala	Gly	Gly	Val	Thr	
	305					310					315					320	
5	Ala	Ala	Ala	Phe	Ala	Val	Gly	Cys	Tyr	Ala	Glu	Trp	Ser	Ser	Val	Ile	
					325					330					335		
	Ile	Ala	Gly	Leu	Thr	Ala	Leu	Met	Gly	Ser	Ala	Phe	Tyr	Tyr	Leu	Phe	
10				340					345					350			
	Val	Val	Tyr	Leu	Gly	Pro	Val	Ser	Ala	Ala	Ala	Ile	Ala	Ala	Thr	Ala	
			355					360					365				
15	Val	Gly	Phe	Thr	Gly	Gly	Leu	Leu	Ala	Arg	Arg	Phe	Leu	Ile	Pro	Pro	
	370						375					380					
	Leu	Ile	Val	Ala	Ile	Ala	Gly	Ile	Thr	Pro	Met	Leu	Pro	Gly	Leu	Ala	
20	385					390					395					400	
	Ile	Tyr	Arg	Gly	Met	Tyr	Ala	Thr	Leu	Asn	Asp	Gln	Thr	Leu	Met	Gly	
					405					410					415		
	Phe	Thr	Asn	Ile	Ala	Val	Ala	Leu	Ala	Thr	Ala	Ser	Ser	Leu	Ala	Ala	
25				420					425					430			
	Gly	Val	Val	Leu	Gly	Glu	Trp	Ile	Ala	Arg	Arg	Leu	Arg	Arg	Pro	Pro	
			435					440					445				
30	Arg	Phe	Asn	Pro	Tyr	Arg	Ala	Phe	Thr	Lys	Ala	Asn	Glu	Phe	Ser	Phe	
	450						455					460					
	Gln	Glu	Glu	Ala	Glu	Gln	Asn	Gln	Arg	Arg	Gln	Arg	Lys	Arg	Pro	Lys	
	465					470					475				480		
35	Thr	Asn	Gln	Arg	Phe	Gly	Asn	Lys	Arg								
					485												



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 01/03980

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12N15/70 C12N15/77 C12P13/08 C12N1/21 //C12N15/31

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C12N C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, SEQUENCE SEARCH, WPI Data, PAJ, MEDLINE, CAB Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	EP 1 085 091 A (DEGUSSA ;KERNFORSCHUNGSANLAGE JUELICH (DE)) 21 March 2001 (2001-03-21) the whole document ---	1-6, 10-13, 16-18
A	WO 98 04715 A (ARCHER DANIELS MIDLAND CO) 5 February 1998 (1998-02-05) the whole document ---	
A	WO 99 53035 A (ALTMAN ELLIOT ;GOKARN RAVI R (US); EITEMAN MARK A (US); UNIV GEORG) 21 October 1999 (1999-10-21) the whole document --- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*G\* document member of the same patent family

Date of the actual completion of the international search

2 October 2001

Date of mailing of the international search report

10/10/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Blanco Urgoiti, B

## INTERNATIONAL SEARCH REPORT

Int. Application No  
PCT/EP 01/03980

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MIZUKAMI T ET AL: "IMPROVEMENT OF THE STABILITY OF RECOMBINANT PLASMIDS CARRYING THE THREONINE OPERON IN AN L-THREONINE-HYPERPRODUCING STRAIN OF ESCHERICHIA COLI W" AGRICULTURAL AND BIOLOGICAL CHEMISTRY, JAPAN SOC. FOR BIOSCIENCE, BIOTECHNOLOGY AND AGROCHEM. TOKYO, JP, vol. 50, no. 4, 1986, pages 1019-1028, XP002047573 ISSN: 0002-1369 the whole document	
A	EIKMANNS B J ET AL: "MOLECULAR ASPECTS OF LYSINE, THREONINE, AND ISOLEUCINE BIOSYNTHESIS IN CORYNEBACTERIUM GLUTAMICUM" ANTONIE VAN-LEEUVENHOEK, DORDRECHT, NL, vol. 64, no. 2, 1993, pages 145-163, XP000918559 the whole document	
A	KRAMER R: "Genetic and physiological approaches for the production of amino acids" JOURNAL OF BIOTECHNOLOGY, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 45, no. 1, 12 February 1996 (1996-02-12), pages 1-21, XP004036833 ISSN: 0168-1656 the whole document	

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/03980

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 1085091	A	21-03-2001	DE 19941478 A1	08-03-2001
			CN 1291651 A	18-04-2001
			EP 1085091 A1	21-03-2001
			JP 2001095592 A	10-04-2001
WO 9804715	A	05-02-1998	AU 730102 B2	22-02-2001
			AU 3899497 A	20-02-1998
			BR 9710503 A	11-01-2000
			CN 1226931 A	25-08-1999
			EP 0917578 A1	26-05-1999
			HU 9903856 A2	28-03-2000
			JP 2000515763 T	28-11-2000
			NO 990362 A	26-01-1999
			PL 331351 A1	05-07-1999
			TR 9900213 T2	21-04-1999
			WO 9804715 A1	05-02-1998
			US 5939307 A	17-08-1999
WO 9953035	A	21-10-1999	AU 3555999 A	01-11-1999
			BR 9909615 A	12-12-2000
			EP 1073722 A1	07-02-2001
			WO 9953035 A1	21-10-1999